

Interpolation & Polynomial Approximation

Divided Differences: A Brief Introduction

Numerical Analysis (9th Edition)

R L Burden & J D Faires

Outline

1 Introduction to Divided Differences

Outline

- 1 Introduction to Divided Differences
- 2 The Divided Difference Notation

Outline

- 1 Introduction to Divided Differences
- 2 The Divided Difference Notation
- 3 Newton's Divided Difference Interpolating Polynomial

Outline

- 1 Introduction to Divided Differences
- 2 The Divided Difference Notation
- 3 Newton's Divided Difference Interpolating Polynomial

Introduction to Divided Differences

A new algebraic representation for $P_n(x)$

Introduction to Divided Differences

A new algebraic representation for $P_n(x)$

- Suppose that $P_n(x)$ is the n th Lagrange polynomial that agrees with the function f at the distinct numbers $x_0, x_1, \dots, x_n$.

Introduction to Divided Differences

A new algebraic representation for $P_n(x)$

- Suppose that $P_n(x)$ is the n th Lagrange polynomial that agrees with the function f at the distinct numbers $x_0, x_1, \dots, x_n$.
- Although this polynomial is unique, there are alternate algebraic representations that are useful in certain situations.

Introduction to Divided Differences

A new algebraic representation for $P_n(x)$

- Suppose that $P_n(x)$ is the n th Lagrange polynomial that agrees with the function f at the distinct numbers $x_0, x_1, \dots, x_n$.
- Although this polynomial is unique, there are alternate algebraic representations that are useful in certain situations.
- The divided differences of f with respect to $x_0, x_1, \dots, x_n$ are used to express $P_n(x)$ in the form

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

for appropriate constants $a_0, a_1, \dots, a_n$.

Introduction to Divided Differences

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

- To determine the first of these constants, a_0 , note that if $P_n(x)$ is written in the form of the above equation, then evaluating $P_n(x)$ at x_0 leaves only the constant term a_0 ; that is,

$$a_0 = P_n(x_0) = f(x_0)$$

Introduction to Divided Differences

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

- To determine the first of these constants, a_0 , note that if $P_n(x)$ is written in the form of the above equation, then evaluating $P_n(x)$ at x_0 leaves only the constant term a_0 ; that is,

$$a_0 = P_n(x_0) = f(x_0)$$

- Similarly, when $P(x)$ is evaluated at x_1 , the only nonzero terms in the evaluation of $P_n(x_1)$ are the constant and linear terms,

$$f(x_0) + a_1(x_1 - x_0) = P_n(x_1) = f(x_1)$$

Introduction to Divided Differences

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

- To determine the first of these constants, a_0 , note that if $P_n(x)$ is written in the form of the above equation, then evaluating $P_n(x)$ at x_0 leaves only the constant term a_0 ; that is,

$$a_0 = P_n(x_0) = f(x_0)$$

- Similarly, when $P(x)$ is evaluated at x_1 , the only nonzero terms in the evaluation of $P_n(x_1)$ are the constant and linear terms,

$$f(x_0) + a_1(x_1 - x_0) = P_n(x_1) = f(x_1)$$

$$\Rightarrow a_1 = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$$

Outline

- 1 Introduction to Divided Differences
- 2 The Divided Difference Notation**
- 3 Newton's Divided Difference Interpolating Polynomial

The Divided Difference Notation

- We now introduce the divided-difference notation, which is related to Aitken's Δ^2 notation [▶ \$\Delta\$ Definition](#)

The Divided Difference Notation

- We now introduce the divided-difference notation, which is related to Aitken's Δ^2 notation [▶ \$\Delta\$ Definition](#)
- The **zeroth divided difference** of the function f with respect to x_i , denoted $f[x_i]$, is simply the value of f at x_i :

$$f[x_i] = f(x_i)$$

The Divided Difference Notation

- We now introduce the divided-difference notation, which is related to Aitken's Δ^2 notation [▶ \$\Delta\$ Definition](#)
- The **zeroth divided difference** of the function f with respect to x_i , denoted $f[x_i]$, is simply the value of f at x_i :

$$f[x_i] = f(x_i)$$

- The remaining divided differences are defined recursively.

The Divided Difference Notation

- The **first divided difference** of f with respect to x_i and x_{i+1} is denoted $f[x_i, x_{i+1}]$ and defined as

$$f[x_i, x_{i+1}] = \frac{f[x_{i+1}] - f[x_i]}{x_{i+1} - x_i}$$

The Divided Difference Notation

- The **first divided difference** of f with respect to x_i and x_{i+1} is denoted $f[x_i, x_{i+1}]$ and defined as

$$f[x_i, x_{i+1}] = \frac{f[x_{i+1}] - f[x_i]}{x_{i+1} - x_i}$$

- The **second divided difference**, $f[x_i, x_{i+1}, x_{i+2}]$, is defined as

$$f[x_i, x_{i+1}, x_{i+2}] = \frac{f[x_{i+1}, x_{i+2}] - f[x_i, x_{i+1}]}{x_{i+2} - x_i}$$

The Divided Difference Notation

- Similarly, after the $(k - 1)$ st divided differences,

$$f[x_i, x_{i+1}, x_{i+2}, \dots, x_{i+k-1}] \quad \text{and} \quad f[x_{i+1}, x_{i+2}, \dots, x_{i+k-1}, x_{i+k}]$$

have been determined,

The Divided Difference Notation

- Similarly, after the $(k - 1)$ st divided differences,

$$f[x_i, x_{i+1}, x_{i+2}, \dots, x_{i+k-1}] \quad \text{and} \quad f[x_{i+1}, x_{i+2}, \dots, x_{i+k-1}, x_{i+k}]$$

have been determined, the **k th divided difference** relative to $x_i, x_{i+1}, x_{i+2}, \dots, x_{i+k}$ is

$$\begin{aligned} & f[x_i, x_{i+1}, \dots, x_{i+k-1}, x_{i+k}] \\ &= \frac{f[x_{i+1}, x_{i+2}, \dots, x_{i+k}] - f[x_i, x_{i+1}, \dots, x_{i+k-1}]}{x_{i+k} - x_i} \end{aligned}$$

The Divided Difference Notation

- Similarly, after the $(k - 1)$ st divided differences,

$$f[x_i, x_{i+1}, x_{i+2}, \dots, x_{i+k-1}] \quad \text{and} \quad f[x_{i+1}, x_{i+2}, \dots, x_{i+k-1}, x_{i+k}]$$

have been determined, the **k th divided difference** relative to $x_i, x_{i+1}, x_{i+2}, \dots, x_{i+k}$ is

$$\begin{aligned} & f[x_i, x_{i+1}, \dots, x_{i+k-1}, x_{i+k}] \\ &= \frac{f[x_{i+1}, x_{i+2}, \dots, x_{i+k}] - f[x_i, x_{i+1}, \dots, x_{i+k-1}]}{x_{i+k} - x_i} \end{aligned}$$

- The process ends with the single **n th divided difference**,

$$f[x_0, x_1, \dots, x_n] = \frac{f[x_1, x_2, \dots, x_n] - f[x_0, x_1, \dots, x_{n-1}]}{x_n - x_0}$$

Generating the Divided Difference Table

x	$f(x)$	First divided differences	Second divided differences	Third divided differences
x_0	$f[x_0]$			
		$f[x_0, x_1] = \frac{f[x_1] - f[x_0]}{x_1 - x_0}$		
x_1	$f[x_1]$		$f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0}$	
		$f[x_1, x_2] = \frac{f[x_2] - f[x_1]}{x_2 - x_1}$		$f[x_0, x_1, x_2, x_3] = \frac{f[x_1, x_2, x_3] - f[x_0, x_1, x_2]}{x_3 - x_0}$
x_2	$f[x_2]$		$f[x_1, x_2, x_3] = \frac{f[x_2, x_3] - f[x_1, x_2]}{x_3 - x_1}$	
		$f[x_2, x_3] = \frac{f[x_3] - f[x_2]}{x_3 - x_2}$		$f[x_1, x_2, x_3, x_4] = \frac{f[x_2, x_3, x_4] - f[x_1, x_2, x_3]}{x_4 - x_1}$
x_3	$f[x_3]$		$f[x_2, x_3, x_4] = \frac{f[x_3, x_4] - f[x_2, x_3]}{x_4 - x_2}$	
		$f[x_3, x_4] = \frac{f[x_4] - f[x_3]}{x_4 - x_3}$		$f[x_2, x_3, x_4, x_5] = \frac{f[x_3, x_4, x_5] - f[x_2, x_3, x_4]}{x_5 - x_2}$
x_4	$f[x_4]$		$f[x_3, x_4, x_5] = \frac{f[x_4, x_5] - f[x_3, x_4]}{x_5 - x_3}$	
		$f[x_4, x_5] = \frac{f[x_5] - f[x_4]}{x_5 - x_4}$		
x_5	$f[x_5]$			

Outline

- 1 Introduction to Divided Differences
- 2 The Divided Difference Notation
- 3 Newton's Divided Difference Interpolating Polynomial

Newton's Divided Difference Interpolating Polynomial

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

Using the Divided Difference Notation

Newton's Divided Difference Interpolating Polynomial

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

Using the Divided Difference Notation

- Returning to the interpolating polynomial, we can now use the divided difference notation to write:

$$a_0 = f(x_0) = f[x_0]$$

Newton's Divided Difference Interpolating Polynomial

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

Using the Divided Difference Notation

- Returning to the interpolating polynomial, we can now use the divided difference notation to write:

$$\begin{aligned} a_0 &= f(x_0) &= f[x_0] \\ a_1 &= \frac{f(x_1) - f(x_0)}{x_1 - x_0} &= f[x_0, x_1] \end{aligned}$$

Newton's Divided Difference Interpolating Polynomial

$$P_n(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)(x - x_1) + \cdots + a_n(x - x_0) \cdots (x - x_{n-1})$$

Using the Divided Difference Notation

- Returning to the interpolating polynomial, we can now use the divided difference notation to write:

$$\begin{aligned} a_0 &= f(x_0) &= f[x_0] \\ a_1 &= \frac{f(x_1) - f(x_0)}{x_1 - x_0} &= f[x_0, x_1] \end{aligned}$$

- Hence, the interpolating polynomial is

$$\begin{aligned} P_n(x) &= f[x_0] + f[x_0, x_1](x - x_0) + a_2(x - x_0)(x - x_1) \\ &\quad + \cdots + a_n(x - x_0)(x - x_1) \cdots (x - x_{n-1}) \end{aligned}$$

Newton's Divided Difference Interpolating Polynomial

$$P_n(x) = f[x_0] + f[x_0, x_1](x - x_0) + a_2(x - x_0)(x - x_1) \\ + \cdots + a_n(x - x_0)(x - x_1) \cdots (x - x_{n-1}).$$

- As might be expected from the evaluation of a_0 and a_1 , the required constants are

$$a_k = f[x_0, x_1, x_2, \dots, x_k],$$

for each $k = 0, 1, \dots, n$.

Newton's Divided Difference Interpolating Polynomial

$$P_n(x) = f[x_0] + f[x_0, x_1](x - x_0) + a_2(x - x_0)(x - x_1) \\ + \cdots + a_n(x - x_0)(x - x_1) \cdots (x - x_{n-1}).$$

- As might be expected from the evaluation of a_0 and a_1 , the required constants are

$$a_k = f[x_0, x_1, x_2, \dots, x_k],$$

for each $k = 0, 1, \dots, n$.

- So $P_n(x)$ can be rewritten in a form called Newton's Divided-Difference:

$$P_n(x) = f[x_0] + \sum_{k=1}^n f[x_0, x_1, \dots, x_k](x - x_0) \cdots (x - x_{k-1})$$

Questions?

Reference Material

Forward Difference Operator Δ

For a given sequence $\{p_n\}_{n=0}^{\infty}$, the **forward difference** Δp_n (read “delta p_n ”) is defined by

$$\Delta p_n = p_{n+1} - p_n, \quad \text{for } n \geq 0.$$

Higher powers of the operator Δ are defined recursively by

$$\Delta^k p_n = \Delta(\Delta^{k-1} p_n), \quad \text{for } k \geq 2.$$

[◀ Return to the Divided Difference Notation](#)